#### (12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

# (19) World Intellectual Property Organization

International Bureau



## T BERTA BEKERNET IN BEGERFRENNE BOEKE BETEL BEGEL EK DIT BOEKE DELDE HERDE BEKEL BETEL ER ET BEFERE BOEK KAN HE

(43) International Publication Date 28 October 2004 (28.10.2004)

PCT

(10) International Publication Number WO 2004/091588 A1

- (51) International Patent Classification<sup>7</sup>: 31/195, 31/198
- A61K 9/50,
- (21) International Application Number:

PCT/EP2004/003669

- (22) International Filing Date:
- 6 April 2004 (06.04.2004)
- (25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

MI2003A000827

18 April 2003 (18.04.2003) I

- (71) Applicant (for all designated States except US): UNI-HART CORPORATION [IE/IE]; 41 Central Chambers, Dublin 2 (IE).
- (72) Inventor; and
- (75) Inventor/Applicant (for US only): TAGLIAMONTE, Alessandro [IT/IT]; Località S. Margherita (La Suvera) 7, I-53019 Castelnuovo Berardenga (IT).
- (74) Agents: BANFI, Paolo et al.; Bianchetti Bracco Minoja S.r.l., Via Plino, 63, I-20129 Milano (IT).

- (81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.
- (84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

#### Published:

- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

WO 2004/091588

PCT/EP2004/003669

# PHARMACEUTICAL COMPOSITION CONTAINING LEVODOPA AND CARBIDOPA

The present invention relates to a prolonged-release oral solid pharmaceutical composition containing the combination levodopa/carbidopa, and the use thereof in the therapy of Parkinson's disease or related pathologies.

#### TECHNICAL BACKGROUND

Parkinson's disease is a neurodegenerative disease that involves different areas of the brain, especially the "substantia nigra", utilizing dopamine as neurotransmitter, causing a slow, progressive disorder of the movements. The main symptoms are bradykinesia, muscular rigidity, resting tremor and postural instability. The diagnosis is usually confirmed by a favourable response to the pharmacological treatment.

The pharmacological therapy is based on the use of selegilins, anticholinergics, amantadines, dopaminergic agonists, ergot alkaloids, levodopa, COMT inhibitors.

Levodopa, upon oral administration, passes the blood-brain barrier and is enzymatically converted to dopamine at the cerebral level.

The choice anti-Parkinson medicament is Sinemet, which contains a combination of levodopa and carbidopa; the latter does not cross the bloodbrain barrier, thus reducing the conversion of levodopa to dopamine by peripheral enzymes, thereby increasing the amount of active ingredient available in the central nervous system.

The therapeutic treatment is usually started with Sinemet 25 mg/100 mg (carbidopa/levodopa) 1/2 tablet a day and after some weeks the dosage is increased until the clinically effective dosage, which is usually 1 tablet 3 or 4 times a day, is reached. Alternatively, treatment with Sinemet CR (sinemet

controlled-release) 50 mg/200 mg starts with 1/2 tablet once a day, and slowly increases until 1 tablet twice a day. The bioavailability of Sinemet CR is approx. 30% lower than that of Sinemet.

Levodopa attains the maximum plasmatic concentration in 1-2 hours and its half-life is 1-3 hours. Therefore, the medicament has to be administered repeatedly during the day and the consequent peaks (Cmax) of plasmatic concentration cause undesired side-effects in the patient. Commercially available carbidopa/levodopa CR (controlled-release) tablets are therapeutically effective over a period of 8 hours, but have various drawbacks and undesired effects, including nausea and vomit, orthostatic hypotension, movement swings, dyskinesias and psychosis.

#### DISCLOSURE OF THE INVENTION

The present invention provides a solid oral pharmaceutical composition containing a combination of carbidopa and levodopa, which ensures a steady release of the active ingredient over 24 hours, thus avoiding plasmatic peaks or fluctuations.

The composition according to the invention contains carbidopa and separately levodopa granules coated by an ethylcellulose film, in a 1:4 carbidopa/levodopa weight ratio. The granular mixture is distributed in a suitable pharmaceutical form, preferably in pre-dosed sachets or hard-gelatin capsules.

The dose of active ingredient per unitary dosage can range from 10 to 200 mg for carbidopa and 40 to 800 mg for levodopa, maintaining a 1:4 weight ratio. The daily dosage can be varied depending on the severity of the disease, the general conditions of the patient, and other parameters. Preferably a 250 mg dose, corresponding to 50 mg carbidopa +200 mg levodopa per unitary dosage, is administered once a day.

According to a preferred embodiment, coated granules are prepared

#### from a mixture containing:

Active ingredient (levodopa or carbidopa)	90%
Polyethylene glycol (carbowax)	3%
Ethylcellulose	1.70%
Talc	0.80%
Polyvinylpyrrolidone	4.50%
Potassium metabisulfite	q.s. (<0.01%)

The granulation process comprises mixing the active ingredient with the binders in a solvent, subsequently granulating the mixture, for example by means of a sieve with suitable mesh to obtain the desired particle size distribution, and coating the granules with ethylcellulose.

According to a preferred embodiment, the granules are prepared by separately mixing the active principles with polyethylene glycol and polyvinylpyrrolidone, and granulating the resulting mixture through a sieve, optionally repeating the process through finer sieves. The coating solution, consisting of ethylcellulose and potassium metabisulfite in acetone and denaturated alcohol, is subsequently sprayed onto the granules, with addition of talc to promote flowing of the granulate; finally coated granules are dried to remove any traces of the solvent. After completing the granulation and coating, coated granules are worked up to the final pharmaceutical form, for example distributed in capsules or sachets.

A once a day administration of the pharmaceutical composition according to the invention is particularly advantageous, in that

- 1) prevents plasma concentration peaks,
- 2) promotes gradual intestinal absorption,
- ensures a steady supply of the active principles, according to predetermined amounts and ratios, during 24 hours.

The therapeutic efficacy and the patient's compliance are therefore

improved, while minimizing the undesired effects.

400 Kg preparation

Thanks to its pharmacokinetic characteristics, the pharmaceutical composition of the invention is conveniently used for the preventive or therapeutical treatment of Parkinson's disease and related disorders.

The following examples illustrate the invention in greater detail.

EXAMPLE 1 - Preparation of the bulk LEVODOPA/CARBIDOPA

400 Kg preparation	
LEVODOPA	359.960 Kg
Carbowax 4000	12.000 Kg
Ethylcellulose	6.800 Kg
Talc	3.200 Kg
Polyvinylpyrrolidone	22.000 Kg
Potassium metabisulfite	0.040 Kg
Acetone*	
Denaturated alcohol*	
Demineralised water*	
100 Kg preparation	
CARBIDOPA	89.990 Kg
Carbowax 4000	3.000 Kg
Ethylcellulose	1.700 Kg
Talc	0.800 Kg
Polyvinylpyrrolidone	4.500 Kg
Potassium metabisulfite	0.010 Kg
Acetone*	
Denaturated alcohol*	

Demineralised water\*

\*solvents used in the process that evaporated off at the end of the process.

#### Preparation of the binder solution

Carbowax 4000 is placed in a stainless steel container fitted with pneumatic stirrer, then polyvinylpyrrolidone is poured, in small amounts, stirring until solubilization.

#### Granulation

Levodopa and Carbidopa are accurately weighed and passed in the granulator using the above binder solution as aggregant.

The wet granulate is forced through a sieve with 840 micron mesh, dried at 40°C for 15 hours in forced-air thermostatised drier and subsequently sieved through a 500 and 840 micron mesh sieve. Powder and granules smaller than 500 micron are regranulated with the same procedure as described above, using deionized water as aggregant. After completion of the granulation process, granules are sieved through a 500 and 840 micron sieve.

The resulting granulate (core) is weighed and placed in a stainless steel basket of the coating pan. During the rotation of the coating pan, which takes place at a suitable rate to ensure an effective rotation of the mass (12 rpm), the binder solution is sprayed onto the granules through a sprayer, preventing formation of drops, and the residual powder active ingredient is added. Spraying of the binder and addition of the powder are carried out at alternate intervals in order to adhere a thin layer of the powder to the core granules and provide better evaporation of the solvent (water) present in the binder solution, which is removed by aspiration avoiding the formation of bubbles. Then the wet granulate is passed through a 1200 micron sieve and dried at 40°C for 15 hours in a forced air thermostatized dryer. After drying, the granulate is sieved again through a 840 and 1200 micron sieve.

#### Preparation of the coating film (solution)

Acetone and denaturated alcohol are placed in the stainless steel container, then ethylcellulose and potassium metabisulfite are added under

continuous stirring, until complete solubilization.

#### Coating of the granules

The granulate from the above step is placed in a fluidized bed and kept in suspension by a filtered air stream.

The coating solution is sprayed intermittently through a sprayer so as to prevent formation of drops. Tale is added to promote flowing of the granulate mass. After completion of the process, the granulate is forced through a 1200 micron sieve. The coated granulate is dried at 40°C for 15 hours in a forced air thermostatized dryer.

#### Preparation of the bulk

The dry coated granulate is sieved through a 840 and 1200 micron sieve and the product is collected in a polyethylene double bag and placed in a metal container with hermetic sealing.

## EXAMPLE 2 - Pharmaceutical preparation in bulk (capsules)

The two granulates (Carbidopa and Levodopa, respectively) obtained as in the examples 1 and 2 above, are distributed into hard-gelatin capsules, maintaining a 1:4 weight ratio (carbidopa/levodopa).

Encapsulation was carried out with a machine equipped with two loading trays, double feeder, two separate dosers (one for each feeder) and programmed to fill the hemi-capsules with the established amounts of granulates.

For filling 250 mg capsules (50 mg carbidopa + 200 mg levodopa), the granulate dosers are set to weigh about 56.6 mg and 222.3 mg of carbidopa and levodopa, respectively.

#### EXAMPLE 3 - Dissolution test

Six samples of a 250 mg preparation (50 + 200) were tested using a solution at pH 1.1 (artificial gastric juice) in a continuous-flow dissolution chamber (25 ml/min, 37°C). The following percentages of release during the 24 hours were obtained (each value is the mean of 6 measurements):

In vitro release	% released average			
Time	Levodopa	Carbidopa		
03™ hour	22%	26%		
06 <sup>rd</sup> hour	43%	39%		
09 <sup>™</sup> hour	59%	51%		
12 <sup>rd</sup> hour	68%	63%		
24 <sup>rd</sup> hour	90%	89%		

The data in the table clearly show that the release of active principles is maintained constant over the entire period of 24 hours.

EXAMPLE 4 – Bioequivalence study between a formulation of the invention (50 mg carbodopa + 200 mg levodopa - denominated Dopabain) and the commercial product Sinemet CR ® (50 mg carbodopa + 200 mg levodopa), upon single administration

In order to evaluate the plasma levodopa concentration of both formulations after 24 hours a single-dose, randomized, two-period, two-sequence crossover bioavalability study was performed in 10 healthy volunteers.

The levodopa concentration of each subject was monitored up to 72 hours. The results of the study are shown in the Figure.


#### **CLAIMS**

- 1. A solid oral pharmaceutical composition containing carbidopa and levodopa in a 1:4 weight ratio, in the form of ethylcellulose-coated granules.
- 2. A pharmaceutical composition as claimed in claim 1, wherein said granules are included in gelatin capsules or sachets.
- 3. A composition as claimed in claim 1, containing from 10 to 200 mg carbidopa and 40 to 800 mg levodopa.
- 4. A composition as claimed in claim 3, containing 50 mg carbidopa and 200 mg levodopa.
- 5. A composition according to any previous claim, containing 90% levodopa and carbidopa, 3% polyethylene glycol, 1.70% ethylcellulose, 0.8% talc, 4.5% polyvinylpyrrolidone, potassium metabisulfite q.s. to 100.
- 6. A composition as claimed in claim 5, wherein the granules have a mean diameter ranging from 500 to 1200  $\mu m$ .
- 7. The use of carbidopa and levodopa in a 1:4 weight ratio for the preparation of ethylcellulose-coated granules to be used in the treatment of Parkinson's diseases or related pathologies.

#### ABSTRACT

An oral solid pharmaceutical composition for constant and prolonged release of a levodopa/carbidopa combination, for use in the treatment of Parkinson disease.

## FIGURE



DOPABAIN-1 DSINEMET-CR1

#### INTERNATIONAL SEARCH REPORT

International Application No
T/EP2004/003669

A. CL	ASSII	FICATION OF SUBJE	CT MATTER	
IPC	7	A61K9/50	A61K31/195	A61K31/198
	•		,	110211027 230

According to International Patent Classification (IPC) or to both national classification and IPC

#### B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols) IPC 7 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ, MEDLINE, EMBASE, BIOSIS

Category °	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,X	WO 03/101432 A (HSIAO CHARLES; HSU LARRY (US); IMPAX LAB INC (US); HAN CHIEN-HSUAN (U) 11 December 2003 (2003-12-11) page 15, lines 1-7 page 16, lines 5-18	1-7
Х	US 2003/031707 A1 (RUBIN ALAN A) 13 February 2003 (2003-02-13) example 3	1-7
x	EP 0 260 236 A (LEJUS MEDICAL AB) 16 March 1988 (1988-03-16) figures 2,3; examples A,B figure 11; example 3 column 4, line 64 - column 5, line 3 claims 1,5,7,8	1-7
V Fudi	er documents are listed in the continuation of box C.    Y   Patent family members are listed in	

X Patent family members are listed in annex.
<ul> <li>*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</li> <li>*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone</li> <li>*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.</li> <li>*&amp;* document member of the same patent family</li> </ul>
Date of mailing of the international search report  07/10/2004
Authorized officer Vermeulen, S

#### INTERNATIONAL SEARCH REPORT

International Application No T/EP2004/003669

	ation) DOCUMENTS CONSIDERED TO BE RELEVANT		12 7 7 7 7 7 7 7 7 7 7 7 7 7 7 7 7 7 7 7	
Category °	Citation of document, with indication, where appropriate, of the relevant passages		Relevant to claim No.	
A	US 4 832 957 A (DEMPSKI ROBERT E ET AL) 23 May 1989 (1989-05-23) column 3, lines 44-63 examples 1-7		1-7	
A	US 5 445 829 A (PARADISSIS GEORGE N ET AL) 29 August 1995 (1995-08-29) column 1, lines 16-20 column 4, lines 58,59 column 6, lines 23-27 column 7, line 10	_	1-7	
,	<del></del>	. •		
	*			
•				
		•		
				•
				•
			·	
	·			
•	- ,			
	·			

## INTERNATIONAL SEARCH REPORT Information on patent family members

International Application No T/EP2004/003669

Pa	atent document		Publication		Patent family		Publication
cited	d in search report		date <sup>.</sup>		member(s)	!	date
WO	03101432	A <sup>-</sup>	11-12-2003	US	200322404		04-12-2003
				US ·	200322836		11-12-2003
				WO	0310143		11-12 <b>-</b> 2003
				US	2004166159	9 A1	26-08-2004
US	2003031707	A1	13-02-2003	US	623869	9 B1	29-05-2001
EP	0260236	Α	16-03-1988	SE	46094		11-12-1989
	٠.	•	•	AT	5829		15-11 <b>-</b> 1990
			,	CA	129819		31-03-1992
				DD	28610		17-01-1991
				DE DK	376620 44228		20-12-1990
				EP	026023		27-02-1988 16-03-1988
				ES	203622		16-05-1988
				FΙ		2 A ,B,	27-02-1988
	•			HŪ	4443	3 A2	28-03-1988
				JP	606240	5 B	17-08-1994
	•			JP	6313912		10-06-1988
				MX	17291		21-01-1994
	•			NO	873589		29-02-1988
				PT SE	8559 860358		01-09-1987 27-02-1988
				US	498169		01-01-1991
us	4832957	A	23-05-1989	AT	8197	 n T	15-11-1992
	1	•••		AU	267338		15-06-1989
				BG	6120	1 B2	28-02-1997
	,			CA	131860		01-06-1993
				DE	387570		10-12-1992
				DE DK	387570		13-05-1993
				EP	68588 032005		25-07-1989 14-06-1989
				ES.	205269		16-07-1994
				GR	300622		21-06-1993
	•			HK	799		24-01-1997
				IE	6154		16-11-1994
			•	JP	194131		23-06-1995
				JP JP	200020 606783		05-01-1990
				LV		7 A4	31-08-1994 20-04-1996
				MX	17412		21-04-1994
				PT		7 A ,B	29-12-1989
				US	490075	5 A	13-02-1990
				US	498340		08-01-1991
				ZA	880918	9 A ·	30-08-1989
บร	5445829	Α	29-08-1995	US	513397		28-07-1992
				US	512238		16-06-1992
				AT AU	12886 63502		15-10-1995
			•	AU	63502 547039		11-03-1993 08-11-1990
				CA	201603		05-11-1990
				DE	6902287		16-11-1995
				DE	6902287		02-05-1996
				EP	039642	5 A2	07-11-1990
				ES US	208079 529623		16-02-1996
				110		rs &	22-03-1994